

VINNIK, K. (Stalinskaya oblast', g.Zhdanov)

Thoughtful friend of steel workers. Okhr.truda i sots.strakh.
3 no.4:62-63 Ap '60. (MIRA 13:6)
(Zhdanov--Steel industry--Hygienic aspects)

VINNIK, K., uchitel'

The activist group is the support of the savings bank. **Fin.**
SSSR 22 no.4:51-52 Ap '61. (MIRA 14:4)
(Zhdanov—Savings banks)

AKOPOV, A.; TUGARINOV, I.; TIMANOVSKIY, I.; NECHAYEV, M.; SEMENOV, V.;
VINNIK, K.; SOMIN, V.

Let us welcome the 22d Congress of the CPSU with excellent achievements. Fin. SSSR 22 no.10:49-59 0 '61. (MIRA 14:9)

1. Zamestitel' nachal'nika Mosgorfinupravleniya (for Akopov).
 2. Zamestitel' zaveduyushchego Leningradskim oblfinotdelom (for Tugarinov).
 3. Nachal'nik byudzhetnogo upravleniya Ministerstva finansov Kazakhskoy SSR (for Timanovskiy).
 4. Zaveduyushchiy Ul'yanovskim oblfinotdelom (for Nechayev).
 5. Zaveduyushchiy Volgodskim oblfinotdelom (for Semenov).
- (Finance) (Taxation)
(Bezhet'sk District--Insurance)

VINNIK, K.

Summer vacation of the petroleum workers of the Chechen-Ingush
A.S.S.R. Neftianik 9 no.9:26 8 '64 (MIRA 18:2)

VINNIK, K.

"Cooking here is better than at home." Obshchestv.pit. no.5:8
My '59. (MIRA 12:8)
(Makhachkala—Restaurants, lunchrooms, etc.)

VINNIK, K.

Improvement of labor protection and safety engineering.
Neftianik 8 no.6:27 Je '63. (MIRA 16:11)

VINNIK, K. (g.Pyatigorsk)

Culinary expert from Platigorsk. Obshchestv.pit. no.4:23-24
Ap '61. (MIRA 14:3)

(Cookery)

VINNIK, K.

VINNIK, K. (Zhdanov)

Azovstal's fire brigade. Pozh.delo 3 no.10:15 0 '57. (MIRA 10:11)
(Zhdanov--Fire prevention)

VINNIK, K.

Automobile club of the university. Neftianik 7 no.9:29 S '62.
(MIRA 16:7)

(No subject headings)

VINNIK, L. A.

"Lymphography and lymphadenography of the lungs in tuberculosis and collapse therapy."

to be presented at the Radiology Congress, Karlovy Vary, Czechoslovakia, 10-14 June 63

VINNIK, L. A.

Vinnik, L. A. - "The functional status of the physiological system of the connective tissues in lung tuberculosis in adults", Trudy Astrakh. gos. med. in-ta, Vol. IX, 1948, P. 111-15.

SO: U-3042, 11 March 53, (Letopis 'Zhurnal 'nykh Statey, No. 3, 1949).

VINNIK, L.A.

Lymphanography and lymphadenography of the lungs in tuberculosis and collapse therapy. Cesk. rentgen. 18 no.4:248-250
Jl'64

1. Astrachansky lekarsky institut a krajska protituberkulezni
poradna.

VINNIK, L. A.

25942 Vinnik, L. A. Gemodinamichekiye sdvigi pri raneniyakh myagkikh tkaney s. zamedlennoy regeneratsiey. V sb: Problemy vosstanovit. Lecheniya invalidov Otechestv. voyny. Astrakhan', 1948, s. 214-19.

S0: Letopis' Zhurnal Statey, No. 30, Moscow, 1948

VINNIK, L.A., kandidat meditsinskikh nauk; CHERVOVA, V.I.

Functional state of the reticulo-endothelial system in tissue
therapy. Klin.med. 33 no.6:86 Je '55 (MLRA 8:12)

1. Iz kafedry fakul'tetskoy terapii (zav.-prof. D.G.Oystrakh)
Astrakhanskogo meditsinskogo instituta.
(TISSUE EXTRACTS) (RETICULO-ENDOTHELIAL SYSTEM)

VINNIK, L.A., kandidat meditsinskikh nauk (Astrakhan')

Some considerations on M.B.TSuker's article "On diseases of the central nervous system of tubercular etiology." Klin.med. 34 no.9: 88-90 S '56. (MIRA 9:11)

1. Iz fakul'tetskoy terapevticheskoy kliniki (sav. prof. D.G.Oystrakh) Astrakhanskogo meditsinskogo instituta i Astrakhanskogo oblastnogo protivotuberkuleznogo dispansera (glavnyy vrach A.P.Demidova)

(CENTRAL NERVOUS SYSTEM, dis.

caused by tuberc.)

(TUBERCULOSIS, compl.

CNS dis.)

YINNIK, L.A., kandidat meditsinskikh nauk (Astrakhan')

Use of calcium and vitamin D in tuberculosis of the lungs. Klin.
med. 34 no.12:82-84 D '56. (MIRA 10:2)

1. Iz Fakul'tetskoy terapevticheskoy kliniki (sav. - prof. D.G.
Oystrekh) Astrakhanakogo meditsinskogo instituta i Astrakhanakogo
oblastnogo protivotuberkuleznogo dispansera (glavnyy vrach A.P.
Demidova)

(TUBERCULOSIS, PULMONARY, ther.
calcium & vitamin D)

(CALCIUM, ther. use
tuberc., pulm., with vitamin D)

(VITAMIN D, ther. use
tuberc., pulm., with calcium)

VINNIK, L.A., kandidat meditsinskikh nauk.

Clinical evaluation of the hippuric acid liver function test;
clinico-anatomical parallels. Terap. arkh. 38 no.1:63-68 '56

(MLRA 9:6)

1. Iz fakul'tetskoy terapevticheskoy kliniki (zav.-prof. D.G.
Oystrakh) i kafedry patologicheskoy anatomii (zav.-prof. G.G.
Nepryakhin) Astrakhanskogo meditsinskogo instituta.

(LIVER FUNCTION TEST,
hippuric acid (Rus))

(HIPPURATES,
liver funct. test (Rus))

VINNIK, L.A., kandidat meditsinskikh nauk

Pharmacological action on the central nervous system in pulmonary tuberculosis. Vrach.delo no.7:761 J1 '57. (MLRA 10:8)

1. Fakul'tetskaya terapevticheskaya klinika (zav. - prof. D.G. Oystakh) Astrakhanskogo meditsinskogo instituta i Astrakhanskogo oblastnogo protivotuberkuleznogo dispansera
(TUBERCULOSIS) (NARCOTICS)

VINNIK, L.A., dotsent; SUTYRINA, G.V.; BOLDYREVA, A.A.; SHEVCHENKO, A.M.

Growth rate of Mycobacterium tuberculosis and isoniazid
concentration in resected lungs. Prob. tub. no.1:75-78 '65.
(MTRA 18:12)

1. Fakul'tetskaya terapevticheskaya klinika (zav.- prof.
A.M. Nogaller) Astrakhanskogo meditsinskogo instituta i
Astrakhanskiy oblastnoy protivotuberkuleznyy dispanser
(glavnyy vrach A.P. Demidova).

VINNIK, I.F.; SIDOROV, K.A.

Early diagnosis and surgical tactics in cancer of the thyroid gland.
Vop. onk. 11 no.10:95-100 '65.

(MIRA 18:10)

1. Iz kliniki Voenno-morskoy i gospi'tal'noy khirurgii (nachal'nik
prof. Ye.V.Smirnov) i kliniki fakul'tetskoy terapii (nachal'nik -
prof. V.A.Beyyer) Voenno-meditsinskoy ordena Lenina akademii im.
S.M.Kirova.

L 59591-65 EWT(1)/TWA(h) Feb 76
ACCESSION NR: AP5017455

UR/0020/65/162/005/1041/1044

15
10
B

AUTHOR: Vinnik, L. P.

TITLE: The structure of 4-6 second microseisms

SOURCE: AN SSSR. Doklady, v. 162, no. 5, 1965, 1041-1044

TOPIC TAGS: microseism structure, microseism observation, seismography 12

ABSTRACT: Microseisms with periods of 4-6 seconds generate serious difficulties during seismological observations. Earlier (mostly three-station) observations seem to indicate that they are generated by oscillations of large water masses. To check these earlier results, the author utilized a new method of directional reception utilizing a system of 7 stations with vertical seismographs and three-component devices. The field data discussed in the paper were collected during a month of observations (October 1961) in eastern Kazakhstan in the Ust'-Kamenogorsk district. All stations utilized the USF seismographs (eigenperiod of 1.5 sec.) with F 117/3 photoelectric amplifiers and GK-VII galvanometers; the registration was accomplished on the RS-2 instrument with a 240 mm/min. development. At 5 sec. the amplification was equal to

Card 1/2

L 59591-65
ACCESSION NR: AP5017455

20,000. The most striking result was the constant presence of a component with a velocity of 10-20 km/sec. and an energy between 10 and 40% of the total energy of the process. Some 4 km/sec. waves (found often) could be surface waves. All the observed waves carried approximately 50% of the total energy; the rest was contained in the irregular component whose nature is not yet clear. The author believes that all components of the microseismic background have the same origin. "The author thanks V. I. Keylis-Borok for the formulation of the problem, and I. P. Pasechnik, N. M. Pruchkina and K. A. Kossova for help during the investigations." Orig. art. has: 4 formulas and 3 figures.

ASSOCIATION: Institut fiziki Zemli im. O. Yu. Shmidta Akademii nauk SSSR
(Institute of Earth Physics of the Academy of Sciences, SSSR)

SUBMITTED: 18Dec64

ENCL: 00

SUB CODE: ES

NO REF SOV: 001

OTHER: 000

Card

2/2

VINNIK, L.A., dotsent; FILIMONOV, Yu.I.

External respiration under controlled physical stress in patients with pulmonary tuberculosis (veloergometric studies). Probl. tub. 42 no.3:27-34 '64. (MIRA 18:1)

1. Fakul'tetskaya terapevticheskaya klinika (zav. - prof. A.M. Nogaller), kafedra fizicheskogo vospitaniya (zav. - I.A. Kolomeytsev) Astrakhanskogo meditsinskogo instituta i Astrakhanskoy oblastnoy protivotuberkuleznyy dispanser.

VINNIK, L.A.

Healing processes in pulmonary tuberculosis under combined
treatment with pneumothorax and antibacterial preparations.
Probl.tub. 38 no.4:33-41 '60. (MIRA 14:5)
(PNEUMOTHORAX) (TUBERCULOSIS)

VINNIK, L.A. (Astrakhan')

"Some remarks concerning Professor B.L. Mazur's article, "Clinical aspects and differential diagnosis of tuberculosis." Kaz. med. zhur. no. 4:100-102 J1-Ag '60. (MIRA 13:8)
(TUBERCULOSIS) (MAZUR, B.L.)

VINNIK, L.A., kand.med.nauk (Astrakhan')

Problem of tuberculosis in general practice. Klin.med. 36
no.12:30-36 D '58. (MIRA 12:6)

1. Iz fakul'tetskoy terapevticheskoy kliniki (zav. - prof.D.G.
Oystrakh) Astrakhanskogo meditsinskogo instituta (dir. - dots.
S.V.Zakharov) i Astrakhanskogo oblastnogo protivotuberkuleznogo
dispansera (glavnyy vrach A.P.Demidova).
(TUBERCULOSIS
in GP (Rus))

VINNIK, L.A., .kand.med,nauk; VISHNEVETSKIY, .F.Ye.; MINSKAYA, .N.M.; PESCHANSKIY, V.S.

Effect of phthivazid on the cardiovascular system in tuberculosis.
Vrach. delo no.1:95-96 '59. (MIRA 12:4)

1. Kafedra fakul'tetskoy terapii (zav. - prof. D.G. Oystakh) i
kafedra patologicheskoy anatomii (zav. - prof. M.S. Brunshteyn)
Astrakhanskogo meditsinskogo instituta.
(ISONICOTINIC ACID) (CARDIOVASCULAR SYSTEM)

VINNIA, L.A., kand.med.nauk

Some problems of therapy in lung abscess. Sov.med. 21 no.8:94-98
Ag '57. (MIRA 10:12)

1. Iz fakul'tetskoy terapevticheskoy kliniki (zav. - prof. D.G.
Oystrakh) Astrakhanskogo meditsinskogo instituta.
(LUNG DISEASES, ther.
abscess (Rus))
(LUNGS, abscess,
ther. (Rus))

VINNIK, L.A.

Therapeutic pneumothorax and antibacterial preparation in complex therapy of pulmonary tuberculosis. Suvrem med., Sofia no.10:64-71 '60.

1. Iz Fakultetskata terapevtichna klinika pri Astrakhanskiia meditsinski institut (Zavezhdasht prof. D.I.Oistrakh [deceased]) i Astrakhanskiia oblasten protivotuberkulozen dispanser (Glaven lekar A.P.Demidova)
(PNEUMOTHORAX ARTIFICIAL)
(ANTITUBERCULAR AGENTS ther)

VINNIK, L.M.; GRINBERG, R.Ya.; KAMINSKIY, Ya.A.; KLEPIKOV, V.D.; KUZNETSOV, A.M.; KUCHENEV, N.I.; STRUZHESTRAKH, Ye.I.; TISHIN, S.D.; KHARITONOV, A.B.; TSEYTS, I.E.; SHAPIRO, I.I.; SHAPIRO, M.Ya.; ANAN'YAN, V.A., retsenzent; VASIL'YEV, D.T., retsenzent; GORETSKAYA, Z.D., retsenzent; KARTSEV, S.P., retsenzent; KEDROV, S.M., retsenzent; KOMISSARZHEVSKAYA, V.N., retsenzent; KOPERBAKH, B.L., retsenzent; KORBOV, M.M., retsenzent; LEONOV, N.I., retsenzent; LUR'YE, G.B., retsenzent; NOVIKOV, V.F., retsenzent; KHARITONOV, A.B., retsenzent; GAL'TSOV, A.D., red.; VOL'SKIY, V.S., red.; KHISIN, R.I., red.; SEMENOVA, M.M., red. izd-va; MODEL', B.I., tekhn. red.

[Handbook for establishing norms in the manufacture of machinery] Spravochnik normirovshchika-mashinostroitelia v tomakh. Moskva, Gos. nauchno-tekhn. izd-vo mashinostroit. lit-ry. Vol.2. [Technical norms for machine-tool operations] Tekhnicheskoe normirovanie stanochnykh rabot. 1961. 892 p. (MIRA 14:11)

(Machinery industry)

(Factory management)

MIKHAYLOV, D.V.; VINNIK, L.M.; SLUCHAYEV, P.N.; SULYAGIN, V.I.;
BARYKOVA, G.I., red.izd-va; GORDEYEVA, L.P., tekhn.red.

[Norms for the wear, strength and consumption of metal-cutting tools] Normy iznosa, stoikosti i raskhoda rezhushchego instrumenta. Moskva, Gos.nauchno-tekhn.izd-vo mashinostroitel'noy lit-ry, 1961. 174 p.

(MIRA 15:2)

1. Russia (1923- U.S.S.R.) Glavnoye upravleniye nauchno-issledovatel'skikh i proyektnykh organizatsiy. Nauchno-issledovatel'skoye byuro tekhnicheskikh normativov. 2. Nauchno-issledovatel'skoye byuro tekhnicheskikh normativov (for Mikhaylov, Vinnik, Sluchayev, Sulyagin).
(Metal-cutting tools--Standards)

VINNIK, L.M.; GRINBERG, R.Ya.; KAMINSKIY, Ya.A.; KLEPIKOV, V.D.; KUZNETSOV, A.M.; KUCHENEV, N.I.; STRUZHESTRAKH, Ye.I.; TISHIN, S.D.; KHARITONOV, A.B.; TSEYTS, I.E.; SHAPIRO, I.I.; SHAPIRO, M.Ya.; ANAN'YAN, V.A., retsenzent; VASIL'YEV, D.T., retsenzent; GORETSKAYA, Z.D., retsenzent; KARTSEV, S.P., retsenzent; KEDROV, S.M., retsenzent; KOMISSARZHEVSKAYA, V.N., retsenzent; KOPERBAKH, B.L., retsenzent; KORBOV, M.M., retsenzent; LEONOV, N.I., retsenzent; LUR'YE, G.B., retsenzent; NOVIKOV, V.F., retsenzent; GAL'TSOV, A.D., red.; VOL'SKIY, V.S., red.; KHISIN, R.I., red.; SEMENOVA, M.M., red. izd-va; MODEL', B.I., tekhn.red.

[Reference book for establishing norms in the manufacture of machinery; in 4 volumes] Spravochnik normirovshchika-mashinostroitelia; v 4 tomakh. Moskva, Gos. nauchno-tekhn. izd-vo mashinostroit. lit-ry. Vol.2. [Establishing technical norms for operating machine tools] Tekhnicheskoe normirovanie stanochnykh rabot. Pod red. E.I.Struzhestrakha. 1961. 392 p.

(MIRA 14:8)

(Industrial management) (Machine tools)

29575
S/049/61/000/005/002/013
D218/D306

9.9865(1109,1327)

AUTHOR: Vinnik, L. P.

TITLE: On grouping low-frequency seismographs

PERIODICAL: Akademiya nauk SSSR. Izvestiya. Seriya geofizicheskaya, no. 5, 1961, 643-648

TEXT: The equivalent circuit of a single seismograph with galvanometer detection is illustrated in Fig. 1. R_s is the resistance of the pendulum coil, r_g is the resistance of the galvanometer frame, R_1 , R_2 and r are additional resistances included in the circuit. The present paper is concerned with a seismograph group connected as shown in Fig. 2 (series-parallel connection), where $\bar{R}_s = R_s + R_1$ and $\bar{r}_g = r_g + R_2$. It is shown that the differential equation relating to the displacement of the soil x , the deflection of the galvanometer φ and the time t is of the form $\ddot{\varphi} + a\dot{\varphi} + b\ddot{\varphi} + c\dot{\varphi} + d\varphi = -e\ddot{x}$ where the coefficients a , b , c , d and e are

Card 1/3

On grouping low-frequency ...

29575
S/049/61/000/005/002/013
D218/D306

constants. It is pointed out that the structure of this differential equation is very similar to the well-known differential equation for a single seismograph; the only difference lies in the values of the constants. A formula is derived for the magnification of a group. The theory is then applied to a one-dimensional group in which the individual seismographs are identical mechanically but have different electrodynamic constants. The frequency characteristics are then given for two-dimensional groups, in particular, a single ring, two rings and four rings. It is suggested that these circular groups may be convenient in detecting first arrivals of body waves due to earthquakes and explosions. It is shown that circular groups can be designed to have highly selective properties in the frequency range 1 - 10 c/s. There are 4 figures and 3 Soviet-bloc references. ✓

ASSOCIATION: Akademiya nauk SSSR, Institut fiziki zemli (Academy of Sciences USSR, Institute of Physics of the Earth)

SUBMITTED: November 15, 1960
Card 2/3

On grouping low-frequency ...

29575
S/049/61/000/005/002/013
D218/D306




Fig. 1




Fig. 2

✓

Card 3/3

L 10742-63

ACCESSION NR: AP3002026

EWI(1)/BDS--AFFTC/ESD-3--TF

8/0049/63/000/006/0350/0860

56

53

AUTHOR: Vinnik, L. P.

TITLE: Space-time filtration of a seismic signal

SOURCE: AN SSSR. Izv. Ser. geofizicheskaya, no. 6, 1963, 850-860

TOPIC TAGS: microseismic noise, grouping of seismographs

ABSTRACT: A mathematical method is outlined for determining the optimum grouping of seismographs on the basis of the computation of the spectral density of interferences. The usual computation of spectral density as a Fourier transform of the three-dimensional function of the intercorrelation of microseismic oscillations in time is too cumbersome when applied to experimental measurements. The proposed method presents several simplified constructions which give satisfactory results in certain cases. Thus, when the microseismic field is isotropic, area study can be replaced by profile study. The computed correlation interval between two points represents the optimum distance between seismographs. When the microseismic background consists of a large number of plane waves propagating with the same velocity in different directions

Card 1/3

L 10742-63

ACCESSION NR: AP3002026

and when the $f_i(t)$ of each wave has the same autocorrelation function $R(\tau)$, the microseismic field is considered to be isotropic. For an individual wave

$$R_1(\tau, r) = R(\tau + r \sin \varphi_1 / v), \quad (1)$$

where v is wave velocity, φ is the angle between the wave front and the direction of propagation along distance r . The intercorrelation of two different waves at any distance will equal zero. The total activity of all waves is expressed as

$$R(\tau, r) = \sum_i R_i(\tau, r). \quad (2)$$

The function $R(\tau)$ can be expanded into the sum of cosines of component harmonics. For $R_1(\tau, r)$, each of these components (if its amplitude is considered to be 1) can be represented by

$$R_1(\tau, r) = \cos(\omega \tau + \omega r \sin \varphi_1 / v). \quad (3)$$

Card 2/3

L 10742-63

ACCESSION NR: AP3002026

3

The harmonic component of this frequency when $R(\tau, r)$ is found by integration at the angle of approach, φ is

$$\frac{1}{2\pi} \int_0^{2\pi} \cos(\omega\tau + \omega r \sin \varphi / v) d\varphi = \cos \omega\tau I_0(\omega r / v), (4)$$

where I_0 is the Bessel zero-order function of the first kind. "The author thanks V. I. Kaylis-Borok and V. F. Pisarenko for discussion and valuable advice." Orig. art. has: 7 figures and 21 formulas.

ASSOCIATION: Akademiya nauk SSSR. Institut fiziki Zemli (Academy of Sciences SSSR. Institute of Physics of the Earth)

SUBMITTED: 07May62

DATE ACQ: 16Jul63

ENCL: 00

SUB CODE: 00

NO REF SOV: 006

OTHER: 001

gcm/1/2
Card 3/3

ACCESSION NR: AP4038146

S/0049/64/000/005/0688/0700

AUTHORS: Vinnik, L. P.; Pruchkina, N. M.

TITLE: Investigating the structure of short period microseisms

SOURCE: AN SSSR. Izv. Seriya geofizicheskaya, no. 5, 1964, 688-700

TOPIC TAGS: microseism, seismograph, random process, surface wave, short period vibration, photoamplifier F 117/3, galvanometer GB IV, oscillograph POB 12

ABSTRACT: Microseisms in the frequency range 0.5-5 cycles have been investigated. The method of analysis is based on the concept of a microseismic background as a statistical homogeneous random process, permitting use of the theory of steady random functions. The model considers two contributing sources: 1) the wave (regular) component and 2) vibrations not propagated to any great distance (irregular). The first is rather strong, the second weak. Data were obtained from three vertical seismographs (natural period of 1.5 sec), with F-117/3 photomultiplier and GB-IV galvanometer attached. The record was obtained on the photographic paper of a POB-12 oscillograph. Field observations were made in eastern Kazakhstan in the summer of 1961, in a hilly area on exposed granite bedrock, far from man-made noise. Correlation analysis indicates that the field of

Cord 1/2

ACCESSION NR: AP4038146

the vortical component in short-period microseisms may have been formed by superposition of two or three surface waves from distant sources (distances greater than the wave length) with irregular vibration. The velocity of the surface waves is computed to be about 3000 m/sec, and the energy contribution is generally more than half the total energy involved in the record. The correlation interval of the irregular vibration is less than 400-600 m. Spectral analysis confirms the view that low-frequency contribution has added to the wave component. Conclusions cannot be final; they need critical analysis by confrontation with new experimental data. It appears probable, however, that the short-period waves are genetically related to longer-period first-order microseisms. The experiments were performed by L. P. Vinnik, programming was attended to by N. M. Pruchkina, field work and data processing were done by A. S. Deniskuv, V. P. Kirillina, and V. N. Tyulyukova. "In conclusion, the authors express their sincere thanks to V. F. Pisarenko, V. I. Keylis-Borok, and I. P. Pasechnik for valuable suggestions and for aid in the work." Orig. art. has: 6 figures, 1 table, and 14 formulas.

ASSOCIATION: Akademiya nauk SSSR Institut fiziki Zemli (Academy of Sciences SSSR, Institute of Physics of the Earth)

SUBMITTED: 26Jun63

DATE ACQ: 12Jun64

ENCL: 00

SUB CODE: ES

NO REF SOV: 002

OTHER: 002

Card 2/2

L 24797-66 EWT(d)/EWP(c)/EWP(v)/T/EWP(k)/EWP(1)/ETC(m)-6 IJP(c)

ACC NR: AP6006952

(N)

SOURCE CODE: UR/0381/65/000/006/0009/0015

AUTHORS: Braynín, E. I.; Vinnikov, L. Ya.

ORG: Institute Giproniselektroshakht, Donetsk (Institut "Giproniselektroshakht")

TITLE: Modeling problem in metal defectoscopy on an EGDA integrator by the potential drop method

SOURCE: Defektoskopiya, no. 6, 1965, 9-15

TOPIC TAGS: / electric potential, defectoscope, electrode potential, model, steel, metal film / EGDA-9-60 integrator

ABSTRACT: A planar modeling technique was used on an EGDA-9/60 integrator to solve the complex potential drop problem in the determination of defects in metals. Two electrode contact methods were used: a defectoscope with electrodes soldered to the metallic surface at two points; and a defectoscope with a thin metallized film deposit adhering to a thick metallic wall. In the first case, equipotential lines are drawn on the metal surface between the two electrodes with and without defects, and the presence and magnitude of the defects are determined from the ratio of potential drop K, with and without the defect. Similar equipotential lines are drawn between the electrodes of the second defectoscope. To evaluate the sensitivity of the method, the potential drop ratio K is plotted against the

UDC: 620.179.18

Card 1/2

L 24797-66

ACC NR: AP6006952

ratio s_1/L , where s_1 is the distance between adjoining electrode current and potential, and L is the overall distance between the electrodes. The sensitivity is seen to increase as L decreases and s_1 increases. Orig. art. has: 7 figures and 1 table.

SUB CODE: 14, 09/ SUBM DATE: 10Jun65/ ORIG REF: 004

Card 2/2

.82

VINNIK, M.A.; AGRANOVSKAYA, A.I.; SEMENOVA, N.N.

X-ray diffraction and microstructure study of phase relations in
the formation of barium-cobalt hexaferrite $\text{Ba}_3\text{Co}_2\text{Fe}_{24}\text{O}_{41}$ (Co_2Z).
Izv.AN SSSR.Neorg.mat. 1 no.7:1177-1183 J1 '65. (MIRA 18:9)

VINNIK, M.A.

Phase relations in the system $BaO - CoO - Fe_2O_3$. Zhur. neorg. khim.
10 no.9:2137-2144 S '65. (MIRA 18:10)

L 21221-66 EWT(m)/EWP(t) LJP(c) JD
ACC NR: AP6003811 SOURCE CODE: UR/0181/66/008/001/0269/0272

AUTHORS: Vinnik, M. A.; Erastova, A. P.; Saksonov, Yu. G.

ORG: none

TITLE: Investigation of the cation distribution in barium
hexaferrites 28
B

SOURCE: Fizika tverdogo tela, v. 8, no. 1, 1966, 269-272

TOPIC TAGS: barrium compound, ferrite, magnetic moment, crystal symmetry, cation, x ray analysis, saturation magnetization

ABSTRACT: The investigated substance $Ba_2Me_2^{2+}Fe_{12}^{3+}O_{22}$ (Me^{2+} stands for Co, Ni, Zn, Cu, Mg, and other metals) has exhibited in various experiments a magnetic moment which differs appreciably from those calculated theoretically by various authors under the assumption that the Me^{2+} ions are located in the spinel blocks of the ferrite structure (E. W. Gorter, Proc. IEE, 104B, Suppl. No. 5, 255, 1957). It is shown, however, that the calculated values and the experimental values

Card 1/2

L 21221-66

ACC NR: AP6003311

of the magnetic moment are in better agreement if it is assumed that Gorter's hypothesis is in error and that the Me^{2+} and Fe^{3+} are statistically distributed over the S and T blocks in both tetrahedral and octahedral positions. The cation distribution in the barium hexaferrite, obtained from the measurements of the specific saturation magnetization by the Weis method (P. Weis, Arch. sci. phys. nat. v. 29, 175, 1910) is then compared with the cation distribution obtained by x-ray structure analysis and the two are shown to be in agreement within the limits of experimental accuracy. The formulas for the cation distributions are tabulated for these ferrites with Mg, Zn, Co, Cu, Mn, and Ni. The dependence of the results on the quenching and cooling of the ferrite is briefly discussed. The data show in addition that in hexaferrites, as in spinels, the Ni^{2+} and Co^{2+} have an affinity to octahedra, while the ions Zn^{2+} and Mn^{2+} to tetrahedra. Orig. art. has: 2 formulas and 2 tables.

SUB CODE: 20/ SUBM DATE: 30Jul65/ ORIG REF: 002/ OTH REF: 004

Card 2/2 *dis.*

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001859830008-9

WINNIE, M.A.

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001859830008-9"

VINNIK, M.A.

Biological accumulation of trace elements in soils under the
forest canopy. Trudy Vor. gos. zap. no.13:179-181 '61.
(MIRA 16:8)
(Voronezh Preserve--Minerals in soil)

J

COUNTRY : USSR

CATEGORY : Soil Science. Physical and Chemical Properties of Soil.

ABS. JOUR. : RZhBiol., No. 4, 1959, No. 15368

AUTHOR : Vinnik, M.A.; Kennova, Ye.K.

INST. : Moscow Univ.

TITLE : Iodometric Determination of Iron in Soils by the Moore-Treadwell Method.

ORIG. PUB. : Vestn. Mosk. un-ta, 1956, No.1, 89-93

ABSTRACT : The Gan-Vindish method of iodometric determination of Fe in soils is widely used in soil-chemical laboratories. The results of determinations by this method are compared to those obtained by the Treadwell modification. The recovered data are identical, but the Moore-Treadwell technique is simpler and permits faster results.

-- F.I. Shcherbak

Card: 1/1

13

L 2301-66

ACCESSION NR: AP5022272

UR/0363/65/001/007/1177/1183
549.73:539.24

AUTHOR: Vinnik, M. A.; Agranovskaya, A. I.; Semenova, N. N.

TITLE: X-ray diffraction and microstructural study of phase relationships in the formation of barium cobalt hexaferrite Ba sub 3 Co sub 2 Fe sub 24 O sub 41 (Co sub 2 Z)*

SOURCE: AN SSSR. Izvestiya. Neorganicheskiye materialy, v. 1, no. 7, 1965, 1177-1183.

TOPIC TAGS: barium compound, cobalt compound, iron compound

ABSTRACT: The object of the work was to study the phase relationships during the formation of Co_2Z and to establish the temperature region of its existence. The compound was synthesized from ferric oxide, cobalt oxide, and barium carbonate by pressing and sintering powder mixtures, and the phase composition of the products was determined by X-ray diffraction and microstructural examination. It is found that Co_2Z does not form directly from the original oxides, but by means of the intermediate compounds $\text{BaFe}_{12}\text{O}_{19}$ (M)* and $\text{Ba}_2\text{Co}_2\text{Fe}_{12}\text{O}_{22}$ (Co_2Y)*. The compound Co_2Z starts to form at 1150C, and is stable when heated in air up to

Card 1/2

L-2301-66

ACCESSION NR: AP5022272

1350C. Above 1350C, it decomposes into $\text{BaCo}_2\text{Fe}_{16}\text{O}_{27}$ (Co_2W), BaFe_2O_4 (B), $\text{BaFe}_{12}\text{O}_{19}$ (M), and the solid solution $\text{Co}_8\text{Fe}_{1-8}\text{O}_4$ (S'), this decomposition is due to the reduction of Fe^{3+} to Fe^{2+} at high temperatures. The compounds Co_2Y and Co_2W also decompose above 1250 and 1300C, respectively. "The authors are deeply grateful to A. A. Shvarts for constant interest in this work." Orig. art. has: 6 figures and 1 table.

ASSOCIATION: None

SUBMITTED: 22Mar65

ENCL: 00

SUB CODE: IC, GC

NO REF SOV: 001

OTHER: 005

*Arbitrary Symbols used by authors to designate various phases in their paper.

M= $\text{BaFe}_{12}\text{O}_{19}$, B= BaFe_2O_4 , S= CoFe_2O_4 , $\text{S}' = \text{Co}_8\text{Fe}_{1-8}\text{O}_4$, $\text{Co}_2\text{Y} = \text{Ba}_2\text{Co}_2\text{Fe}_{12}\text{O}_{22}$,

$\text{Co}_2\text{Z} = \text{Ba}_2\text{Co}_2\text{Fe}_{24}\text{O}_{41}$, $\text{Co}_2\text{W} = \text{BaCo}_2\text{Fe}_{16}\text{O}_{27}$

Card

2/2

ACC NR: AP7006228

(A)

SOURCE CODE: UR/0078/67/012/001/0038/0043

AUTHOR: Vinnik, M. A.; Agranovskaya, A. I.; Semanova, N. N.

ORG: none

TITLE: X-ray diffraction and microstructural study of the phase relationships in the formation of barium cobalt hexaferrite $\text{BaCo}_2\text{Fe}_{16}\text{O}_{27}$

SOURCE: Zhurnal neorganicheskoy khimii, v. 12, no. 1, 1967, 38-43

TOPIC TAGS: barium compound, cobalt compound, ferrite

ABSTRACT: The compound $\text{BaCo}_2\text{Fe}_{16}\text{O}_{27}$, designated Co_2W , was synthesized by a solid phase reaction from Fe_2O_3 , Co_3O_4 and BaCO_3 , and the phase composition of the products (kept at 500-1440°C for 4-20 hr, then quenched in air or furnace-cooled) was determined by x-ray diffraction and microstructural methods. Co_2W does not form directly from the initial oxides, but via the intermediates $\text{BaFe}_{12}\text{O}_{19}$ (barium hexaferrite) and cobalt ferrite CoFe_2O_4 . The compound Co_2W begins to form at 1150°C and is stable at temperatures below 1350°C. At 1350°C and above, Co_2W decomposes, the products being $\text{BaFe}_{12}\text{O}_{19}$ and $\text{Co}_8\text{Fe}_{11}\text{Fe}_2^{3+}\text{O}_4$. The decomposition is due to the reduction of Fe^{3+} to Fe^{2+} at high temperatures. Orig. art. has: 7 figures and 1 table.

SUB CODE: 07/ SUBM DATE: 26Feb65/ ORIG REF: 004/ OTH REF: 001

Card 1/1

UDC: 546.732'723'431-31:539.26

VINNIK, M.I.
CA

*Heterogeneous acid-base catalysis in multimolecular
Alms. N. M. Chirkov and M. I. Vinnik. Doklady Akad.
Nauk S.S.S.R. 88, 1007-1010 (1947). Transformation of
AcII into parahexyle was studied between 60° and 100°. When completely pure, parahexyle does not decomp. into
AcII at these temps. Treatment of app. walls with HCl
vapor catalyzes the decomp., and the rate rises sharply
with increase of temp. (results given graphically). When
 P/P_{decomp} is over 0.4, the reaction-rate const. rises rapidly
with increased vapor pressure of parahexyle: $K \propto$
 $K_{\text{ad}}/P_{\text{a}}$. Up to P/P_{a} 0.3 the content of HCl goes linearly
with the amt. of adsorbed HCl, but above 0.3 the linearity
does not hold, and it is estd. that in the expts. up to 60 mol.
layers of HCl were adsorbed on the walls, at P/P_{a} of 0.75.
At P/P_{a} 0.5-0.6 the activation energy of depolymerization
is 26 kcal./mole. The catalytic activity of the wall-
adsorbed HCl drops with introduction of H_2O vapor; a small
decline occurs with EtOH vapors. Apparently in these
cases HCl dissociates and does not participate in the catalysis.
G. M. Kosolapoff*

VINNIK, M. I.

USSR/Physical Chemistry - Polymerization of butylene on phosphoric acid.

Card 1/1

Authors : Chirkov, N. M. and Vinnik, M. I.

Title : Kinetics and mechanism of reactions in the presence of thin films of non-volatile acids.

Periodical : Dokl. AN SSSR 95, 6, 1243 - 1247, 21 Apr 1954

Abstracts : Thin films of non-volatile acids coating hard bodies behave as continuous heterogenous catalysts. The article gives partial data on an experiment performed in a laboratory and describes conditions under which the process mentioned can be observed. Diagrams.

Institution : Institute of Chemical Physics of the Acad. of Scs. of the USSR

Submitted : 13 Feb, 1954

USSR/Chemistry - Physical chemistry

Card 1/1 Pub. 22 - 41/56

Authors : Churkov, N. M., and Vinnik, M. I.

Title : Kinetics and mechanism of reactions in the presence of thin films of non-volatile acids. The kinetic equation for isobutylene isomerization over H_3PO_4

Periodical : Dok. AN SSSR 99/5, 823-826, Dec 11, 1954

Abstract : The kinetic laws governing the process of isobutylene polymerization at 40 - 90° temperatures and 40 - 600 mm mercury column pressures in the presence of an acid (H_3PO_4) catalyst are explained. A kinetic equation is presented which makes it possible to describe not only the initial rates of polymerization but also the kinetic curves provided certain constants formulated for different initial isobutylene pressures are used at a given temperature and acid concentration. The independence of the rate of reaction from the pressure of the reagents in the case of a heterogeneous film catalyst was found to be the result of a maximum possible absorption of the olefine by the acid. One USSR reference (1954). Table; graphs.

Institution: Academy of Sciences USSR, Institute of Chemical Physics

Presented by: Academician V. N. Kondratyev, June 29, 1954

Vinnik, M.I.

Math ✓ Kinetics and mechanism of reactions catalysed by thin films of non-volatile acids: polymerization of isobutylene in contact with aqueous sulphuric acid. N. M. Chirkov and M. I. Vinnik (*Dokl. Akad. Nauk, SSSR*, 1955, 105, 766-769).—Experiments were carried out on the polymerization of isobutylene in vessels packed with quartz or glass tubes moistened with aq. H_2SO_4 of various concn. There was a linear relation $\log K_1 + 1.7 H_0 = \text{const.}$, between the bimolecular velocity const. K_1 and the acidity function H_0 of the acid film. The plot of $\sqrt{v_i}/P$ against $\sqrt{v_i}$, where v_i is the initial reaction velocity (in mm./min.) and P is the isobutylene pressure (in mm.) is linear at a given temp. and acid concn. A general expression is obtained for the initial reaction rate with 70% H_2SO_4 . F. W. KIRKBRIDE.

2 .
MA-YOUT2
JCCP'CS

5002

PM

Inst. Chem. Phys., AS USSR

VINNIK, M. I.

✓ 12216* (Russian.) Problem of Preparing Pure Boron Fluoride. K voprosu o poluchenii chistogo storistogo bora. M. I. Vinnik, G. B. Manelis, R. S. Riabova, G. D. Tantsyrev, and N. M. Chirkov. *Zhurnal Neorganicheskoi Khimii*, v. 1, no. 4, 1956, p. 628-631.
Method of preparing KBF_4 and preparing and purifying BF_3 .
Mass spectral and infra-red analyses.

Vinnik K, M. I.

USSR/Physical Chemistry - Solutions. Theory of Acids and Bases, B-11

Abst Journal: Referat Zhur - Khimiya, No 1, 1957, 489

Author: Vinnik, M. I., Kruglov, R. N., and Chirkov, N. M.

Institution: None

Title: Acidity of Aqueous Solutions of Hydrobromic and Hydrochloric Acid

Original

Periodical: Zh. fiz. khimii, 1956, Vol 30, No 4, 827-836

Abstract: The indicator method was used in measuring the acidity H_0 of aqueous solutions of HBr(I) and HCl(II) over the concentration ranges 0.275-56.52 wt percent and 8.9-40.47 wt percent, respectively. From the experimental values of H_0 , values were calculated for $(f_{H_2O} + f_B)f_{BH} +$ (III) and $f_B/(f_A - f_{BH})$ (IV). The standard state is chosen such that the acid ionization constant $[HA/K_0] = 1$. It is shown that the ratio III increases with increasing concentrations of I and II and the ratio IV is practically independent of the concentrations of I and II and is equal to one. For aqueous solutions of I and II up to 16-17 ml, the acidity H_0 is numerically equal to $\log(a_{HA}/c_A)$, where a_{HA} is the

Card 1/2

USSR/Physical Chemistry - Solutions. Theory of Acids and Bases, B-11

Abst Journal: Referat Zhur - Khimiya, No 1, 1957, 489

Abstract: activity of the acid determined from the emf or from the vapor pressure and c_A - is the concentration of the halide ion.

Card 2/2

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001859830008-9

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001859830008-9"

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001859830008-9

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001859830008-9"

20-6-27/47

Vinnik, M. I.

AUTHORS: Vinnik, M. I. , Ryabova, R. S. , Chirkov, N. M.

TITLE: The Kinetics of the Acid-Catalytic Decarbonylation of Benzoyl Formic Acid (Kinetika kislotno-kataliticheskogo dekarbonilirovaniya benzoilmurav'inoi kisloty)

PERIODICAL: Doklady AN SSSR, 1957, Vol. 117, Nr 6, pp. 1017 - 1020 (USSR)

ABSTRACT: The present paper investigates the determination of particles participating in the elementary act of the acid processes and compares the constant of speed with the concentration of the particles. The reactions $C_6H_5COCOOH \rightarrow CO + C_6H_5COOH$ were selected as model. The solutions of benzoyl formic acid in water and in diluted solutions of sulphuric acid do not absorb in the visible range of the spectrum. On dissolution of $C_6H_5COCOOH$ in concentrated H_2SO_4 (80 - 90 %) an absorption in the visible range of the spectrum appears. The absorption coefficient of benzoyl formic acid considerably increases with increasing concentration of H_2SO_4 . In the present work the speed of the decarbonylation of solutions of $C_6H_5COCOOH$ in H_2SO_4 was determined from the speed of the decrease in the optical density of the solution when $\lambda = 400 m\mu$. The optical density was measured by the spectrometer ($\Phi - 4$) in quartz cuvettes. A diagram illustrates a typical kinetic curve of the decarbonylation

Card 1/3

20-6-27/47

The Kinetics of the Acid-Catalytic Decarbonylation of Benzoyl Formic Acid

process (the dependence of the optical density D of the solution on the time t) and its logarithmic anamorphosis. The influence of the acidity h upon the constant K of the speed was investigated in the range of concentrations of from 85,46 to 99,94 % of H_2SO_4 at $T = 15^\circ C$. The corresponding results are illustrated by a diagram. In the range from $h_0 = 10^{8,06}$ to $h_0 = 10^{9,4}$ the dependence between $lg K$ and the acidity function H_0 is linear: $lg K + 1,8 H_0 = \text{const}$. At higher values of acidity a deviation from the linearity was observed. The temperature dependence of K for the solutions of $C_6H_5COCOOH$ in 98,80; 96,2; 92,89,77 and 85,46 % H_2SO_4 were determined in a narrow range of temperature. These data are in agreement with the Arrhenius (Arrhenius) equation. Benzoyl formic acid is supposed to exist in different forms in the concentrated and diluted solutions of H_2SO_4 . In diluted H_2SO_4 it is supposed to exist in a nonionized state, but with increasing acidity of the medium it is ionized. Then the authors report on the determination of the constant of alkalinity. There are 4 figures, 1 table, and 1 non-Slavic reference.

Card 2/3

^{20-6-27/47}
The Kinetics of the Acid-Catalytic Decarbonylation of Benzoyl Formic Acid

ASSOCIATION: Institute of Chemical Physics AS USSR
(Institut khimicheskoy fiziki Akademii nauk SSSR)

PRESENTED: June 17, 1957, by V. N. Kondrat'yev, Academician

SUBMITTED: June 12, 1957

AVAILABLE: Library of Congress

Card 3/3

20-119-1-26/52

AUTHORS: Vinnik, M. I., Manelis, G. B., Eppele, G. V., Chirkov, N. M.

TITLE: Kinetics of Isobutylene Polymerization in the Presence of Boron Fluoride Etherate (Kinetika polimerizatsii izobutilena v prisutstvi efirata ftoristogo bora)

PERIODICAL: Doklady Akademii Nauk SSSR, 1958, Vol. 119, Nr 1, pp. 98-100 (USSR)

ABSTRACT: The present paper investigates the polymerization of iC_4H_8 in the presence of a complex compound of the diethyl ether with boron fluoride: $(C_2H_5)_2O \cdot BF_3$. The catalyst $(C_2H_5)_2O \cdot BF_3$ in the form of a thin adsorbed film was applied to the surface of little tubes of melted quartz for the purpose of avoiding diffusion-conditioned disturbances. The reaction container with an insertion of little quartz tubes was evacuated to a pressure of from 10^{-4} - 10^{-5} mm of mercury column previous to the experiment. At first the necessary pressure of ether vapor was produced in the reaction container and then the boron fluoride was introduced. In every experiment the pressures of etherate (p_{etherate}^g) of boron fluoride ($p_{BF_3}^g$), of ether in the gaseous phase (p_{ether}^g), corresponding

Card 1/3

20-119-1-26/52

Kinetics of Isobutylene Polymerization in the Presence of Boron Fluoride Etherate

to the equilibrium and the quantity of the etherate ($P_{\text{etherate}}^{\text{fl}}$) condensed on the surface were determined. The data used for the determination of these values are given in brief. Special attention was paid to the production of the pure reagents which must not contain any traces of moisture. The reaction velocity was expressed by the reduction of the isobutylene pressure referring to 1 mole of the adsorbed etherate. A diagram shows the kinetic curve and its anamorphosis for the polymerization process of iC_4H_8 in the presence of the etherate $(C_2H_5)_2O.BF_3$. If $t = 70^\circ C$ and $P_{iC_4H_8}$ is small

(up to 100 - 1500 mm of mercury column), the kinetic equation up to a 40 - 50 per cent transformation can easily be expressed by a secondary order equation. The constant of velocity K_1 thus determined does not depend on the initial pressure of the iC_4H_8 . The influence of ether on the catalytic activity of the $(C_2H_5)_2O.BF_3$ is similar to the influence of water on the acidity of mineral acids (phosphoric acid, sulfuric acid etc.). Such an intense influence of the ether - even at low concentrations - can only be explained by its

Card 2/3

20-119-1-26/52

Kinetics of Isobutylene Polymerization in the Presence of Boron Fluoride Etherate

basic properties. A diagram and a table illustrate the dependence of the constant K_p of the polymerization velocity on the pressure of boron fluoride in gaseous phase corresponding to the equilibrium. The etherate of boron fluoride is an effective catalyst for the polymerization of iC_4H_8 . Judging from the catalytic activity the etherate must have the properties of an intense acid. There are 3 figures, 1 table, and 7 references, 3 of which are Soviet.

ASSOCIATION: Institut khimicheskoy fiziki Akademii nauk SSSR
(Institute for Chemical Physics AS USSR)

PRESENTED: August 7, 1957, by V. N. Kondrat'yev, Member, Academy of Sciences, USSR

SUBMITTED: August 1, 1957

Card 3/3

BEL'SKIY, V.Ye.; VINNIK, M.I.

Kinetic method of the analysis of mixtures of acetic anhydride
and acetic acid. Zhur. anal. khim. 19 no.3:375-378 '64.
(MIRA 17:9)

1. Institut khimicheskoy fiziki AN SSSR, Moskva.

5 (4)

AUTHORS:

Manelis, G. B., Vinnik, M. I.
Chirkov, N. M. (Moscow)

SOV/76-33-5-11/33

TITLE:

The Acid Function of the System Diethyl Ether - BF_3
(Funktsiya kislotnosti sistemy dietilovyy efir - BF_3)

PERIODICAL:

Zhurnal fizicheskoy khimii, 1959, Vol 33, Nr 5,
pp 1030-1034 (USSR)

ABSTRACT:

The acid function was investigated over a wide concentration range by using various indicators: 2,4-dinitroaniline, 2,4-dichloro-6-nitroaniline, 5-Cl-2-nitroaniline, and o-nitroaniline. The measurement was carried out by means of the spectrophotometer SF-4 at room temperature. The compound $(\text{C}_2\text{H}_5)_2\text{OBF}_3$ was synthesized in a quartz cuvette. Table 1

shows the dependence of $\lg \frac{C_B}{C_{\text{BH}^+}}$ on the concentration of the

ether compound for the individual indicators (C_B = concentration of the non-ionized indicator, C_{BH^+} = concentration of the ionized indicator). The determination of the acid function

Card 1/3

The Acid Function of the System Diethyl Ether - BF_3

SOV/76-33-5-11/33

was carried out assuming that the ether compound investigated is a strong acid and that indicators in strong acids have the same value of pK_B (negative logarithm of the basicity constant). Therefore, $\text{pK}_B = -4.4$ was put for 2,4-dinitroaniline, and pK_B for the other indicators determined by the usual method. The possible inaccuracies of the determination are discussed. Table 2 and a figure give the results of the determination. The reasons for the deviation of the data obtained from the results mentioned in reference 3 could not be found. Starting from the equilibrium constant the equation of the acid function for $(\text{C}_2\text{H}_5)_2\text{OBF}_3$ in $(\text{C}_2\text{H}_5)_2\text{O}$ is deduced:

$$\text{H}_0 = \lg \sqrt{k_1 \left(\frac{100}{[\text{H}]} - 1 \right) + k_2 \cdot (\text{C}_2\text{H}_5)_2\text{OBF}_3}$$
 surpasses the phosphoric acid because of its acidity degree ($\text{H}_0 = -6.22$ at 100°). The quantitative data of the polymerization rate of isobutylene are in good agreement with it. There are 1 figure, 2 tables, and 7 references, 5 of which are Soviet and 1 Czechoslovakian.

Card 2/3

The Acid Function of the System Diethyl Ether - BF_3 SOV/76-33-5-11/33

ASSOCIATION: Akademiya nauk SSSR Institut khimicheskoy fiziki Moskva
(Academy of Sciences of the USSR, Institute of Chemical
Physics, Moscow)

SUBMITTED: October 9, 1957

Card 3/3

SOV/76-33-9-17/37

5(4)

AUTHORS:

Vinnik, M. I., Ryabova, R. S., Chirkov, N. M.

TITLE:

Kinetics and Mechanism of Reactions in Concentrated Strong Acid Media. I. Kinetics of Dehydration of o-Benzoyl Benzoic Acid in Concentrated Solutions of Sulphuric Acid

PERIODICAL:

Zhurnal fizicheskoy khimii, 1959, Vol 33, Nr 9, pp 1992-2001 (USSR)

ABSTRACT:

An investigation was made of the kinetics of catalytic dehydration of o-benzoyl benzoic acid (I) to anthraquinone in sulphuric acid (79.92% to 100%) at temperatures of from 50° to 97.5°C. The spectrophotometrical method (spectrophotometer type SF-4) was applied and the dependence of the dehydration rate constant on the acidity of the medium and on temperature was determined (Table 1). Three forms of (I) were found to occur in the state of equilibrium in the case of (I) being ionized in highly acid medium - a non-ionized form BOH, a protonized form BOH_2^+ , and a dehydrated protonized form B^+ . The ratio between the concentrations of the three forms is determined by the acidity of the medium and the activity of water. The limiting stage of the process is the isomerization of the dehydrated

Card 1/3

SOV/76-33-9-17/37

Kinetics and Mechanism of Reactions in Concentrated Strong Acid Media.

I. Kinetics of Dehydration of o-Benzoyl Benzoic Acid in Concentrated Solutions of Sulphuric Acid

form B⁺. A specification is given of the values of the actual rate constants at 25°C, as obtained by extrapolation (according to the Arrhenius Law), further, the concentration of the three forms of (I) at various sulphuric acid concentrations (Table 2), as well as the dependence of the ionization constant on acidity (Table 3). By spectrometric and kinetic methods the following values were obtained:

$$K_1 = \frac{a_{H^+} a_{BOH}}{a_{BOH_2^+}} = 4 \cdot 10^8; K_2 = \frac{a_{H^+} a_{BOH_2^+}}{a_{B^+} a_{H_2O^+}} = 1 \cdot 10^9$$

Values were obtained for the true activation energy $E_w = 24.5$ kcal/mol (characteristic of the energy balance of the elementary act of the isomerization of dehydrated (I)-ion to anthraquinone), the sum of the ionization heats $Q_1 + Q_2 = 6.5$ kcal/mol, and the value of the true preexponential factor $A = 7 \cdot 10^{11}$ sec⁻¹. There are 7 figures, 3 tables, and

Card 2/3

SOV/76-33-9-17/37

Kinetics and Mechanism of Reactions in Concentrated Strong Acid Media.

I. Kinetics of Dehydration of o-Benzoyl Benzoic Acid in Concentrated Solutions of Sulphuric Acid

8 references, 1 of which is Soviet.

ASSOCIATION: Akademiya nauk SSSR, Institut khimicheskoy fiziki, Moskva
(Academy of Sciences of the USSR, Institute of Chemical
Physics, Moscow)

SUBMITTED: February 22, 1958

Card 3/3

5(4)

AUTHORS:

Utyanskaya, E. Z., Stepanyants, A. U., SOV/20-124-5-38/62
Vinnik, M. I., Chirkov, N. M.

TITLE:

The Calculation of the Function of Acidity and the Molecular Composition of Hydrofluoric Acid From the Data of the Nuclear Magnetic Resonance of F^{19} (Raschet funktsii kislotnosti i molekulyarnogo sostava plavkovoy kisloty po dannym yadernogo magnitnogo rezonansa F^{19})

PERIODICAL:

Doklady Akademii nauk SSSR, 1959, Vol 124, Nr 5, pp 1095-1098 (USSR)

ABSTRACT:

Hammett's function of acidity is measured by means of the indicator method and amounts to $X_0 = -\lg a_{H^+}(f_B/f_{BH^+})$. Here f_B and f_{BH^+} denote the coefficients of the activities of the ionized and not ionized forms of the indicator; a_{H^+} - the proton activity in the given medium. In the aqueous solutions of HF there are the following kinds of equilibrium: $HF \rightleftharpoons H^+ + F^-$ (2); $HF + F^- \rightleftharpoons HF_2^-$ (3); $K_1 = a_{H^+}a_{F^-}/a_{HF}$ (4); $K_2 = a_{HF_2^-}/a_{HF}a_{F^-}$ (5).

Card 1/3

The Calculation of the Function of Acidity and the
Molecular Composition of Hydrofluoric Acid From the Data of the Nuclear
Magnetic Resonance of F^{19} SOV/20-124-5-38/62

Here $K_1 = 6.89 \cdot 10^{-4}$ and $K_2 = 2.695$ denote the constants of equilibrium, expressed in activities. From the dissociation equation (4) of hydrofluoric acid there follows:
 $-\lg a_{H^+} = -\lg K_1 + \lg (a_{F^-}/a_{HF})$. If the quantities K_1 , a_{HF} and m_{F^-} are known, it is possible to calculate the acidity function H_0 . For the purpose of determining quantitative results concerning the composition and the acidity of concentrated aqueous solutions of HF the authors carried out measurements of the chemical shifts of the resonance of F^{19} in aqueous solutions of hydrofluoric acid of different concentrations. For the chemical shift to be observed and also for a system consisting of several interacting components expressions are written down. The chemical shift to be observed is of the order of magnitude $(5.95 \text{ to } 6.05) \cdot 10^{-4}$ for the various concentrations of hydrofluoric acid. In order, therefore to be able to determine the acidity function with an accuracy

Card 2/3

The Calculation of the Function of Acidity and the
Molecular Composition of Hydrofluoric Acid From the Data of the Nuclear
Magnetic Resonance of F^{19} SOV/20.124.5-38/62

of ± 0.05 , it is necessary to know the observable chemical shift with an accuracy of $\pm 0.03 \cdot 10^{-4}$. Measurements were carried out in a magnetic field of ~ 3300 oersted. In the course of these measurements a saw-tooth modulation of the magnetic main field was used. The results obtained by measurements are shown by a table and by a diagram. At concentrations of up to 30 % the variations of the chemical shift are within the limits of measuring errors. The results obtained by calculating molecular composition and the acidity function are shown by tables and diagrams. There are 4 figures, 2 tables, and 9 references, 5 of which are Soviet.

ASSOCIATION: Institut khimicheskoy fiziki Akademii nauk SSSR (Institute of Chemical Physics of the Academy of Sciences, USSR)
PRESENTED: October 29, 1958, by V. N. Kondrat'yev, Academician
SUBMITTED: October 22, 1958
Card 3/3

SOV/20-126-6-43/67

5(4)

AUTHORS:

Vinnik, M. I., Zarakhani, N. G., Medvetskaya, I. M.,
Chirkov, N. M.

TITLE:


The Role of Salt Formation in Acid-catalytic Processes. The
Kinetics of Cyclohexanone Oxime Hydrolysis (O roli soleobrazo-
vaniya v kislотно-kataliticheskikh protsessakh. Kinetika
gidroliza tsiklogeksanonoksima)

PERIODICAL:

Doklady Akademii nauk SSSR, 1959, Vol 126, Nr 6, pp 1300-1303
(USSR)

ABSTRACT:

The hydrolysis of amides and oximes is accelerated by acids
and bases. An anomalous dependence of the reaction rate upon
the acidity of the medium is observed in acid hydrolysis
(Refs 1-4). The value of the reaction constant increases with
increasing HCl-concentration up to a certain amount and then
decreases again. An investigation of the bimolecular reaction
between the "protonized" molecule of the reagent and water :



of hydrolysis shows that this reaction is not able to explain
the phenomena observed. The authors assume that the "protonized"
form of the reagent is able to combine with the acid anion to a

Card 1/2

The Role of Salt Formation in Acid-catalytic Processes. SOV/20-126-5-43/67
The Kinetics of Cyclohexanone Oxime Hydrolysis

non-dissociated salt. In order to prove the rightness of this assumption, the hydrolysis of cyclohexanone oxime was spectrophotometrically investigated at $\lambda = 222 \text{ m}\mu$ under the catalytic effect of hydrochloric acid. A kinetic curve is derived (Fig 1) under consideration of a monomolecular reaction of the hydrolysis of the oxime and (in the case of a reverse process) of the bimolecular oxime formation from cyclohexanone and hydroxylamine. Table 1 and figure 2 show the experimental results and confirm that the reaction constant of hydrolysis passes through a maximum. A decreasing rate of hydrolysis (Table 2) was brought about by experiments made with the addition of NaCl and LiCl. Therefore, salt formation inhibits acid catalytic processes by reducing the concentration of the reactive form of the reagent. In the hydrolysis of cyclohexanone oxime, the stage of isomerization of the "protonized" form has a limiting effect. There are 2 figures, 2 tables, and 7 references, 2 of which are Soviet.

PRESENTED: February 26, 1959, by V. N. Kondrat'yev, Academician

SUBMITTED: February 20, 1959
Card 2/2

PLANE 1 BOOK EXPLORATION NOV/99

Problem listed in bulletin. (v) 10: *Pratica e Teoria-chimica* bulletins (Problems of Kinetics and Catalysis). (vol.) 10: *Propricce and Proprietà* (Kinetics of Catalysis) Moscow, Izdatno AN SSSR, 1960. 461 p. Errata slip inserted. 2,600 copies printed.

Ref.: S.2. Regularly, Corresponding Member of the Academy of Sciences USSR and A.T. Krylov, Candidate of Chemistry; M. of Publishing House: A.T. Shubnikov Tech. Sci. D.A. Asvat'yev.

PURPOSE: This collection of articles is addressed to physicists and chemists and to the community of scientists in general interested in recent research on the physics and physical chemistry of catalysis.

CONTENTS: The articles in this collection were read at the conference on the Physical and Chemical Properties of Catalysts organized by the USSR Academy of Sciences and Russian Academy of Chemical Sciences, Academy of Sciences (USSR) National Scientific Center on the Problem of "the intensive bioengineering trial in catalysis". The Conference was held at the USSR Academy of Sciences (Institute of Physical Chemistry) from April 20-29, 1986.

Of the great volume of material presented at the conference, only papers not published elsewhere were included in this collection.

XX. FOR GENERAL PURPOSES OF CATEGORICAL

214	Smith, A.S. Spectroscopy of Simple Processes on the Surface of Oxide Catalysts
225	Pyridic, W.E. Types of Active Complexes and Their Role in Heterogeneous Catalysis
240	Shilovskiy, L.A. [Acidimetric Institute] [Laboratory Transports/Moscow Institute of Transportation Engineering] Some Problems of Organic Catalysts
	IV. ACID - BASE CATALYSIS
247	Toshimura, K.Y. [Department of Chemistry of Keio University] Studies on the Activity of Amphoteric Catalytic Catalysts
255	Gilley, R.H. [Institute of Chemical Physics of the U.S. USSR] Mechanism of Homogeneous Acid Catalysis and Its Relation to Heterogeneous Acid Catalysis
273	Debye, O.Y. [Institute of Physical Chemistry of the U.S. USSR] Catalysts Acting on Solid Phases
279	Kono, Hideo, Kenji, Kogakurei, and Y. Sakami. [Chemical Institute of the National Academy of Science, Kyoto] Phosphoric Acid Catalyst for the Isomerization of Glycerol
285	Myakis, M.Y., O.S. Maslovskiy, and B.M. Gidlovskiy [Institute of Chemical Physics of the U.S. USSR] Catalytic Properties of the Complex Compounds of Boron Fluoride
291	Pillay, V.R. and D.S. Prasad [Department of Physics of Indian State University] Spectral Investigations of the Action of Various Aqueous Catalysts
292	Schneider, Th. O. [Laboratory of Macromolecular of Industrial State University] Kinetics of Copies and Structures of the Surface of Silica Catalysts
296	Tseng, T.T., S.Y. Dornayevskiy, Yu. S. Izrael, Z.Y. Gershteyn, V.A. Kuznetsov, I.G. Kozlovskiy, and V. V. Shalovskiy [Research Institute of Chemical Engineering] [Kiev] [Soviet Union] Catalytic Research Institute of Feedstock Products and By-products. The Analysis of Reactions Catalyzed by Liquid and Solid Acids
298	Adams, R.C., and G.L. Ingold [Institute of Physical Chemistry of the University] Experimental Investigation of the Correspondence Between Kinetic and Thermodynamic Acid Base Catalysis
301	Reagan, L.J. [MIT] Ion-pairing effect in gas-liquid catalysis
	Transition-metal catalysts catalyze (Aluminum chloride, a powerful catalyst of Petroleum Refining and the Production of Synthetic Liquid Fuel).
	Contributions to the Problem of Selecting Synthetic Stable Oxide Catalytic Catalysts
	December, 1958. (Aluminum chloride, a powerful catalyst of Petroleum Refining and the Production of Synthetic Liquid Fuel). Acid Properties and Catalytic Capacity of Catalysts

VINNIK, M.I.; MANELIS, G.B.; CHIRKOV, N.M.

Catalytic properties of complex compounds of boron fluoride. Probl.
kin. i kat. 10:285-290 '60. (MIRA 14:5)

1. Institut khimicheskoy fiziki AN SSSR.
(Boron fluoride)

VINNIK, M.I., kand.khimicheskikh nauk; GONIKBERG, M.G., doktor khimicheskikh nauk

Mechanism of heterolytic reactions. Vest.AN SSSR 30 no.9:
115-118 S '60. (MIRA 13:9)
(Chemical reactions, Rate of)

VINNIK, M.I.; ZARAKHANI, N.G. (Moscow)

Kinetics and mechanism of reactions in concentrated strong acid media. Part 3: Kinetics of the hydrolysis of cyclohexanone oxime in hydrochloric and sulfuric acid media. Zhur. fiz. khim. 34 no.12: 2671-2681 D '60. (MIRA 14:1)

1. AN SSSR, Institut khimicheskoy fiziki.
(Cyclohexanone) (Acids, Inorganic)

VINNIK, M.I.; MOISEYEV, Yu.V.; PALAGINA, L.V.

Kinetics of caprolactam hydrolysis in aqueous solutions of KOH.
Dokl.AN SSSR 138 no.1:149-152 My-Je '61. (MIRA 14:4)

1. Institut khimicheskoy fiziki AN SSSR. Predstavleno akademikom
V.N.Kondrat'yevym.
(Hexamethylenimine)

VINNIK, M.I.; ZARAKHANI, N.G. (Moscow)

Kinetics and mechanism of reactions in concentrated strong
acid media. Part 8. Zhur. fiz. khim. 38 no.4:907-915 Ap '64,
(MIRA 17:6)

1. Akademiya nauk SSSR, Institut khimicheskoy fiziki.

ZARAKHANI, N.G.; VINNIK, M.I. (Moscow)

Composition and equilibria in aqueous solutions of carboxylic acids. Zhur. fiz. khim. 38 no.3:632-638 Mr '64.

(MIRA 17:7)

1. Institut khimicheskoy fiziki AN SSSR.

VINNIK, M.I.; MOISEYEV, Yu.V.; PALAGINA, I.V.

Kinetics and the mechanism of γ -butyrolactam hydrolysis
in potassium hydroxide solutions. Kin. i kat. 5 no.2:253-262
Mr-Apr '64. (MIRA 17:8)

1. Institut khimicheskoy fiziki AN SSSR.

BEL'SKIY, V.Ye.; IVANKOVA, N.I., VINNIK, M.I.

Kinetics of the acylation of nitroanilines in boron fluoride
solutions in glacial acetic acid. Zhur. fiz. khim. 39
no.6:1426-1431 Je '65. (MIRA 18:11)

1. Institut khimicheskoy fiziki AN SSSR. Submitted March
10, 1964.

LIBROVICH, N.B.; VIKHAR, M.I.

Activity coefficients of the nonionized form of some indicators used in measuring the acidity of aqueous solutions of sulfuric acid. Dokl. AN SSSR 166 no.3:647-650 Ja '66.

(MIRA 19:1)

1. Institut khimicheskoy fiziki AN SSSR. Submitted May 24, 1965.

Zakharov, A., POPOV, V.V., VINNIK, M.I.

Reaction and mechanism of reactions taking place in concentrated strong acid media. Part 10. Zhur. fiz. khim. 39 no.8.1863-1868 Apr '65. (MIRA 18:9)

1. Institut khimicheskoy fiziki AN SSSR.

MOISEYEV, Yu.V.; ORLOV, I.G.; VINNIK, M.I.

Effect of nonelectrolytes on the infrared spectrum of water. Part 1:
Hydration of butyrolactam in aqueous, alkaline, and acid solutions.
Zhur. struk. khim. 6 no.3:387-390 My-Je '65.

(MIRA 18:8)

1. Institut khimicheskoy fiziki AN SSSR.

ZARAKHANI, N.G.; BODYLINA, V.V.; VINNIK, M.I.

Kinetics and mechanism of reactions taking place in a medium of strong concentrated acids. Part 9. Zhur.fiz.khim. 39 no.7:1561-1565 J1 '85.

(MIRA 18:8)

1. Institut khimicheskoy fiziki AN SSSR.

VINNIK, M.I.; BEL'SKIY, V.Ye.; IVANKOVA, N.I.

2,4,6-Trinitroaniline acylation kinetics and the determination of equilibrium concentrations of ions in solutions of boron fluoride in acetic acid. Zhur.fiz.khim. 39 no.7:1624-1630 J1 1965.

(MIRA 18:8)

1. Institut khimicheskoy fiziki AN SSSR.

ACADEMIC PRESS

U.S. DEPARTMENT OF ENERGY
NATIONAL BUREAU OF STANDARDS


AUTHOR: Vinnik, M. I.; Bel'skiy, V. Ye.; Ivankova, N. L.

TITLE: Kinetics of acylation of 2,4,6-trinitroaniline and determination of equilibrium ion concentrations in acetic acid solutions of boron fluoride

SOURCE: Zhurnal fizicheskoy khimii, v. 39, no. 7, 1965, 1624-1630

TOPIC TAGS: acylation, trinitroaniline, trinitroanilide, boron fluoride, acetic acid

ABSTRACT: The acylation of 2,4,6-trinitroaniline in concentrated boron fluoride solutions in glacial acetic acid is a reversible process in which the conversion of 2,4,6-trinitroaniline to the anilide depends on the BF_3 content of the solution. Conversion can be increased by adding acetic anhydride or decreased by adding water. These facts were utilized to calculate the equilibrium concentrations of the solutions containing from 20 to 40% of the anhydride. The following processes:



Card 1/2

I 63467-65

ACCESSION NR: AP5019790

The equilibrium of the acylation process is represented by the equation
 $2,4,6\text{-trinitroaniline} + \text{BF}_3 \cdot \text{CH}_3\text{COOH} \rightleftharpoons 2,4,6\text{-trinitroanilide} + \text{BF}_3 \cdot \text{H}_2\text{O}$. From the dependence of the anilide: aniline equilibrium ratio on the amount of anhydride added, the concentrations of the anilium ions were calculated. As the amount of anhydride increases, the rate constant of anilide formation increases. It was found that the anilide ion concentration increases with the amount of anhydride added. The equilibrium constant of the anilide formation was calculated. It was found that the change in the activity of the anilide ion is proportional to the change in the activity of the anilium ion. The equilibrium constant of the anilide formation was calculated. It was found that the change in the activity of the anilide ion is proportional to the change in the activity of the anilium ion.

ASSOCIATION: Indicator: Aniline. The equilibrium constant of the anilide formation was calculated. It was found that the change in the activity of the anilide ion is proportional to the change in the activity of the anilium ion.

REMITTANCE: Main

IC

Card 2/2

BEL'SKIY, V.Ye.; VINNIK, M.I. (Moscow)

Acid properties of boron fluoride solutions in acetic acid. Zhur.fiz.
khim. 38 no.8:1956-1955 Ag '64. (MIRA 18:1)

1. Institut khimicheskoy fiziki AN SSSR.

MOISEYEV, Yu.V.; BAKHRAKH, E.Ya.; VINNIK, M.I. (Moscow)

Kinetics of hydrolysis of N-methylacetamide in KOH aqueous
solutions. Zhur. fiz. khim. 37 no.4:784-790 Ap '63.
(MIRA 1967)

1. Akademiya nauk SSSR, Institut khimicheskoy fiziki.

VINNIK, M.I.; RYABOVA, R.S. (Moscow)

Use of the indicator method for determining the concentration
of ions and nondissociated particles. Zhur. fiz. khim. 38 no.3:
606-615 Mr '64. (MIRA 17:7)

1. Institut khimicheskoy fiziki AN SSSR.

BEL'SKIY, V.Ye.; VINNIK, M.I.

Kinetics and the mechanism of acylation of aromatic amines in
the system acetic acid - acetic anhydride. Izv.AN SSSR.

Ser.khim. no.1:40-45 Ja '64.

(MIRA 17:4)

1. Institut khimicheskoy fiziki AN SSSR.

ZARAKHANI, N.G.; VINNIK, M.I.

Determination of the dimer concentration in aqueous solutions
of formic acid. Zhur. fiz. khim. 37 no.11:2550-2553 N'63.

(MIRA 17:2)

1. Institut khimicheskoy fiziki AN SSSR.